

## Anticoagulants, Non-Vascular Surgeons

<b>Medications to stop 7 days prior</b>	<b>Antiplatelets:</b> Agrylin (Anagrelide), Plavix (Clopidogrel), Ticlid (Ticlopidine), Dipyridamole (Aggrenox, Persantine), Pentoxifylline (Trental), Cilostazol (Pletal), Effient (Prasugrel), Brilinta (Ticagrelor) <b>All NSAIDS</b>
<b>Medications to stop 5 days prior</b>	<b>Coumadins:</b> Jantoven, Warfarin
<b>Medications to stop 2 days prior</b>	<b>All other anticoagulants:</b> Xarelto (Rivaroxaban), Pradaxa (Dabigatran), Eliquis (Apixaban), Arixtra (Fondaparinux)
<b>Medications to stop 24 hours prior</b>	<b>Heparin Anticoagulants:</b> Lovenox (Enoxaparin, Lmwh), Heparin, Fragmin (Dalteparin)

**Important Notes:** Patient(s) may require a different blood thinner to protect them while off their regular anticoagulant.

## Anticoagulants, Vascular Surgeons

<b>Medications to stop 5 days prior</b>	Warfarin
<b>Medications to stop 2 days prior</b>	Apixaban (Eliquis), Dabigatran Etixilate (Pradaxa), Edoxaban, Rivaroxaban (Xarelto), Clopidogrel (Plavix) <b>Plavix note:</b> Vascular does not usually stop this medication for the following procedures: <i>Angiogram, lower extremity carotid procedures, CEFI, TCAR</i>
<b>Medications to stop 12 hours prior</b>	Enoxaparin (Lovenox) HEPARIN: Individualized; refer to physician instructions