

# Medications to Hold Before Surgery

## Medications to stop

### 14 days prior

Selegiline, Ensam, Eldepryl, Zelapar (MAOI)

## Medications to stop

### 7 days prior

All injectable GLP-1 agonists

## Medications to stop

### 4 days prior

Ertugliflozin (Steglatro, Steglujan)

## Medications to stop

### 3 days prior

Canagliflozin (Invokana, Invokamet)  
Dapagliflozin (Farxiga, Xigduo, Qtern)  
Empagliflozin (Jardiance, Glyxambi, Synjardy)

## Medications to hold day of surgery

All oral agents, short acting  
GLP-1 agnoists and all non-insulin  
therapy. Okay to take  
the day before.

Accupril (ACE)  
Aceon (ACE)  
Actos (Diabetes)  
Agrrenox (ASA = dipyramidole)  
Aldactone (diuretic)  
Altace (ACE)  
Amaryl (Diabetes)  
Ansaid or Flurbiprofen (NSAID)  
Atacand (Angio 2 blocker)  
Avalide (Irbesartan+HCTZ) (Angio 2  
blocker+diuretic)  
Avapro (Angio 2 blocker)

Azor (Olmesartan+Amlodipine) (Angio  
2 Inh+Ca Channel blocker)

Benazepril (ACE)

Bextra (NSAID)

Bumetanide or Bumex (diuretic)

Byetta or Exenatide (Diabetes)

Candesartan or Atacand or Cilexetil  
(Angio 2 blocker)

Captopril (ACE)

Celebrex (NSAID) *(follow-up with  
surgeon with questions)*

Chlorthalidone (diuretic)

Cozaar or Losartan (Angio 2 Inh)

Diabeta (Diabetes)

Enalapril (ACE)

Etodolac (NSAID)

Exforce (Diovan+Norvasc) (Angio 2  
blocker+antianginal)

Feldene or Piroxicam (NSAID)

Fosinopril (ACE)

Glipizide (Diabetes)

HCTZ (diuretic)

Hyzaar (Cozaar+HCTZ) (Angio 2  
inh+diuretic)

Ibuprofen (NSAID)

Indapamide or Lozol (diuretic)

Indomethacin (NSAID)

Ketoprofen (NSAID)

Ketorolac or Toradol (NSAID)

Lasix (diuretic)

Lisinopril (ACE)

Lodine (NSAID)

Lotensine (ACE)

Lotrel (Amlodipine+Benzapril) (Ca Ch  
blocker+ACE)

Lozol (diuretic)

Mavik (ACE)

Maxzide (diuretic)

Meloxicam, Miloxicam, or Mobic  
(NSAID)

Metformin (Diabetic)

Metolazone (diuretic)

Microzide or HCTZ (diuretic)

Monopril (ACE)

Motrin (NSAID)

Mycardis or Telmisartan  
(Angio 2 blocker)

Naprosyn (NSAID)

Olmesartan (Angio 2 Inh)

Piroxicam or Feldene (NSAID)

Plavix (blood thinner)

Pletal (Platelet Inh)

Precose or Acarbose (Diabetes)

Prinivil (ACE)

Quinapril (ACE)

Ramipril (ACE)

Relafen or Nabumetone (NSAID)

Spirolonactone (diuretic)

Sulindac (NSAID)

Ticlid (Platelet Inh)

Toradol (NSAID)

Voltaren (NSAID)

Zaroxolyn or Mykrox (diuretic)

Zestoretic (Lisinopril+HCTZ,  
ACE+diuretic)

Zestril (ACE)

Ziac (Bisoprolol+HCTZ) (beta block,  
diuretic+Antihypertensive)

**IMPORTANT NOTES:** If Hypoglycemic the morning of surgery, drink 4-8 oz of clear apple juice depending on the severity of the hypoglycemia.